

Using a Holistic Approach to Achieve Weight Loss Maintenance:

Results from the Spirited Life Intervention

AUTHORS

Rae Jean Proeschold-Bell, PhD*

Duke Global Health Institute, Duke Center for Health Policy & Inequalities Research
Duke University

Dori Steinberg, PhD, RD

Duke School of Nursing, Duke Global Health Institute
Duke University

Jia Yao, MA and David E. Eagle, PhD

Duke Center for Health Policy & Inequalities Research
Duke University

Timothy W. Smith, PhD

Department of Psychology
The University of Utah

Grace Cai

Trinity College of Arts & Sciences
Duke University

Elizabeth L. Turner, PhD

Department of Biostatistics and Bioinformatics, Duke Global Health Institute
Duke University

Corresponding author:

Rae Jean Proeschold-Bell, PhD
Duke Global Health Institute, Duke Center for Health Policy & Inequalities Research
Box 90392
Durham, NC 27708-0392
Tel: (919) 613-5442
Fax: (919) 613-5466
Rae.jean@duke.edu

Abstract

Weight loss maintenance is essential to sustain the

health benefits of weight loss. Studies with lower intensity intervention supports under real-world conditions are lacking.

Purpose. Examine changes in weight and cardiometabolic biomarkers among Spirited Life participants following initial 12-month weight loss at 12-24 months and 24-42 months.

Methods. 719 clergy received a wellness intervention including a 10-week online weight-loss program in the first 12 months and monthly health coaching throughout 24 months. Mean changes in weight, blood pressure, high-density lipoproteins and triglycerides were estimated using random effects linear models, accounting for repeated measures. Weight was additionally analyzed in subsamples stratified by body mass index (BMI).

Results. At baseline, 17.1% of participants had BMI < 25 kg/m² and 11.8% had BMI ≥ 40 kg/m². Mean 12-month weight loss was -2.4kg (95% CI: -2.8kg, -2.1kg). On average, at 42 months, participants regained weight but did not exceed baseline (-0.5kg, 95% CI: -1.2kg, 0.2kg), improvements in triglycerides were completely sustained (-13.9mg/dL, 95% CI: -18.6 mg/dL, -9.2 mg/dL), and systolic blood pressure improvements remained significant (-1.9mmHg, 95% CI: -3.0 mmHg, -0.9 mmHg). Participants with a BMI ≥ 40 kg/m² lost significantly more weight that was sustained at 42 months (-5.8kg, 95% CI: -8.9kg, -2.7kg).

Conclusions. The Spirited Life wellness intervention produced weight loss and, for participants with higher levels of obesity, sustained weight loss maintenance. The intervention was effective for long-term prevention of weight gain among participants with BMI of 25 to ≤ 40 kg/m², through 42 months. Wellness interventions such as Spirited Life should be considered for adoption.

KEYWORDS

Weight loss maintenance; clinical trial; holistic health; clergy; health behavior change; occupation

This is an author-produced PDF of an article accepted for publication in the *Translational Behavioral Medicine* following peer review. The full article citation is: Proeschold-Bell, R.J., Steinberg, D., Yao, J., Eagle, D.E., Smith, T.W., Cai, G., & Turner, E.L. (2018). Using a holistic health approach to achieve long-term weight maintenance: Results from the Spirited Life intervention. *Translational Behavioral Medicine*, doi: 10.1093/tbm/iby117. The final publication is available at: <https://academic.oup.com/tbm/advance-article-abstract/doi/10.1093/tbm/iby117/5239983?redirectedFrom=fulltext>

Implications

Practice: Long-duration, lower-intensity interventions that use a variety of ways to teach behavior change strategies can support long-term weight maintenance and sustain the associated health benefits.

Policy: Employers who want to prevent weight gain and reduce chronic disease among employees should consider offering weight loss and behavior change programs to all employees regardless of current health status, and engaging employees for relatively long periods of time (1-2 years) through periodic workshops, healthy eating education, health coaching, and small grants.

Research: Future research should determine the cost-benefit of offering weight maintenance programs to all versus some employees and determine the optimum length of intervention to maximize long-term health benefits.

Introduction

Obesity is one of the largest global causes of preventable morbidity and mortality [1]. This risk of morbidity and mortality decreases with weight loss [2]. Standard behavioral weight loss interventions, using lifestyle changes in diet and exercise habits, produce an average loss of 7-10% of initial weight at one year [3-5]. Yet, even small changes in body weight – losses of 3-5% of initial body weight – can produce clinically meaningful improvements in health [6]. Despite the efficacy of behavioral interventions for weight loss, many individuals who lose weight struggle to maintain improvements post-intervention [7]. Maximum weight losses from behavioral interventions commonly occur at around six months, with participants gradually regaining their weight thereafter [8; 9]. If there is no emphasis on maintenance past 6 months, about half of participants will regain back to their starting weight; over a two-year period post-treatment, most individuals regain one-third to two-thirds of the weight they lost, and some gain beyond their baseline weight [10-13]. In the gold standard Look AHEAD trial, even four years after receiving a lifestyle intervention, about 50% of participants who lost between 5-10%

regained close to or above starting weight [14]. It is critically important to identify ways to support people in weight loss maintenance [15].

In addition, weight stability is now considered a health-promoting goal. US adults from age 18-55 typically gain 0.5-1.0 kg per year [16]. While it varies by age, studies on men suggest average weight gains of 0.18-0.29 kg per year [16; 17]. The 2013 obesity treatment guidelines from the American Heart Association, the American Academy of Cardiology, and The Obesity Society not only recommend weight loss for all overweight and obese individuals, but also state that individuals with obese or overweight status who are not interested in weight loss should focus on preventing weight gain [6]. Importantly, weight stability can confer health benefits; some studies have shown that individuals who maintain their weight for upwards of 20 years can improve or keep stable cardiometabolic indicators such as blood pressure, glucose, and lipid levels [18].

Whether it be for weight loss, weight loss maintenance, or weight stability, providing multi-faceted support for lifestyle modification remains one of the most effective elements of a behavioral weight control intervention [19]. Both the weight loss and weight loss maintenance phases of obesity treatment emphasize skill building for behaviors such as portion control, relapse prevention, incorporating activities into one's lifestyle, social support, and accountability. Yet, some strategies do differ when entering the maintenance phase of a behavioral weight control intervention [20]. The National Weight Control Registry indicates that successful maintainers weigh frequently, stay consistent with their eating pattern, and exercise at levels higher than what is typically promoted for weight loss [21]. Efficacy trials testing interventions for weight loss maintenance incorporating these strategies have shown positive results [10; 22]. For example, in the STOP Regain trial, which emphasized daily weighing as the main self-regulation strategy, Wing and colleagues (2006) found that fewer participants who received support via face-to-face meetings regained weight compared to those who received no support; however, support via digital channels was effective at preventing regain as well, albeit less effective compared to face-to-face [22]. Similarly, Svetkey and colleagues (2008) compared an intervention of monthly personal contact versus an interactive technology-based intervention versus a self-directed control arm; both intervention arms focused on motivation, self-monitoring, problem-solving, and support for 30 months [10]. At 30 months post-intervention, the personal-contact group weighed 1.2 kg less than the interactive technology-based group, although even the technology-based group outperformed the self-directed control group at 18 months. These results suggest that intensive support is most effective for preventing regain and maintaining weight loss, however, even minimal support with a lighter touch for weight maintenance can be effective. A lighter touch intervention (e.g., less in-person contact and fewer minutes per month than, for example, the Look AHEAD intervention [14]) can be less burdensome to participants and has greater potential to be implemented in real-world settings [23].

The workplace offers a means of reaching people for weight loss and weight gain prevention interventions, in a setting where people can support each other's efforts [24]. However, it is unclear whether lighter touch workplace interventions can lead to long-term outcomes. In one large-scale trial of hotel employees that provided group sessions (mean attendance=5.4) and supportive messages in the workplace for almost a year, weight loss was

not significantly different at 12 or 24 months [25]. Other lighter touch workplace interventions have similarly produced null effects [26].

We conducted a pragmatic behavioral intervention trial called Spirited Life. It included lighter touch weight intervention components among an occupational group and its goals were to decrease metabolic syndrome, produce weight loss at 12 months among those who were overweight or obese, maintain weight loss through and beyond the two-year intervention period, and prevent weight gain among participants of normal weight. The intervention combined evidence-based components with some innovations (see Figure 1). Participants of all weight statuses were encouraged to sign up for a ten-week, online, weight-loss program called Naturally Slim® that emphasized eating only when hungry; decreasing sugar intake; eating smaller portions; and balancing fats, proteins, and carbohydrates. Naturally Slim® did not have a rigorous evidence base. The company shared with us findings from hospital groups that enrolled highly motivated patients who evidenced weight loss maintenance and metabolic syndrome reduction several years post-intervention based on medical record data, but with no control group or adjustment for loss to follow-up [27]. However, Naturally Slim's nutritional, portion control, and accountability approaches were consistent with weight loss recommendations [19] and had the added advantage of being scalable for a dispersed group. To this online intervention, we offered follow-up support through up-to-monthly phone health coaching sessions. Health coaches utilized motivational interviewing, which has a strong evidence base for a wide range of behavior changes [28-32]. They encouraged participants to set their own health goals, which could involve physical health, mental health, spiritual well-being or all three. For weight loss, health coaches focused on the Naturally Slim weight loss strategies, and for weight loss maintenance they revisited these health strategies plus continued to engage in goal-setting and motivational interviewing techniques. Health coaching has been shown to be effective for weight loss and related health behaviors [33-38].

One way in which we intended the intervention to be innovative was in its long duration of 24 months; we wanted time to allow participants to change behaviors, slip up in the course of normal life fluctuations (e.g., change in seasons, work expectations, family life), and still have support to return to healthy practices [39]. We built in three components to engage participants in such a long intervention that was also sporadic in that participants did not need to set aside regular time, for example in the way weight loss groups require. One component for ongoing engagement was monthly health coaching. Another was punctuating the intervention with in-person workshops which occurred at the beginning, middle, and end of the two years. For this clergy population, each workshop included theological reasons to attend to one's health and were packaged in ways to resonate with clergy (e.g., morning prayer, communion). The first workshop focused on initial motivation to lose weight and make other holistic health behavior changes. The second and third workshops sought to maintain weight loss, with content focused on motivating additional behavior change, promoting health behavior self-efficacy through teaching the process of behavior change, and providing opportunities for participants to articulate core values, re-commit to behavior change, and plan for sustaining their accomplishments. The third component was offering participants \$500 grants one year into the intervention to assist them in meeting health goals and to maintain their motivation; we also considered this a means of promoting weight loss maintenance. At the time, there was not evidence of the efficacy of financial incentives for weight loss

maintenance [40; 41], although anecdotally health insurance companies were considering them.

The Spirited Life intervention had other components less relevant to the current study and described elsewhere [39]. They included stress management content, although there was no evidence in the two-year outcomes that stress symptoms had improved. They also included targeting people who interact with each other and who might provide mutual support or be influenced by a change in social norms. Although we did not measure it specifically, there was anecdotal evidence that clergy talked to each other about weight loss strategies and noticed weight loss in each other. Many participants commented that they felt a shift in climate towards self-care being more acceptable for clergy.

We invited all United Methodist clergy in North Carolina (NC) to participate, irrespective of current health status. We included normal weight clergy consistent with a selective prevention strategy [42]. Christian clergy are in great need of weight loss and metabolic syndrome reduction programs. Across multiple denominations, the prevalence of obesity in US Christian clergy is estimated to be 39% to 43% [43-48]. In a previous study with this population, we found the obesity prevalence was 10.3 percentage points higher than the surrounding comparable population [43]. These same studies found high rates of chronic disease among clergy. In our Spirited Life study, 50.9% met criteria for metabolic syndrome [49].

The Spirited Life study aims were to: 1) for participants overweight or obese at baseline, promote initial weight loss and reversal of metabolic syndrome components (central obesity, elevated triglycerides, low high-density lipoproteins, hypertension, and elevated blood glucose) [50]; 2) for participants overweight or obese at baseline, maintain weight loss and reversal of metabolic syndrome components for at least 18 months post-intervention; and 3) for participants with normal weight at baseline, promote healthy weight stability. The intervention lasted 24 months. We randomly assigned participants to one of three cohorts: immediate intervention, one-year waitlist cohort, and two-year waitlist cohort. After the intervention, we followed two of the three cohorts for an additional 18 months.

After two years, intention-to-treat analyses revealed significant reductions in weight, averaging -1.7 kg to -3.0 kg, depending on intervention cohort [49]. The 505 participants who were obese at baseline evidenced an estimated intervention effect of 1.8 kg (95% CI: 0.01 kg, 3.62 kg) greater weight loss compared to control participants. These levels of weight loss are modest, but consistent with effectiveness trials conducted in real-world settings [51]. In addition to weight loss, the prevalence of metabolic syndrome in Spirited Life was 12% lower at the end of the two-year intervention.

In the current paper, we examine the impact of the Spirited Life intervention on weight loss maintenance after initial weight loss but during low-intensity intervention (months 12-24), and after the intervention (months 24-42), as well as on long-term changes in cardiometabolic biomarkers (e.g., blood pressure) impacted by weight. We further examine long-term weight stability in normal weight participants.

Methods

Procedure

We have described Spirited Life trial methods in a protocol article [39] and reported on two-year outcomes covering the intervention period [49]. Briefly, we contacted all clergy who were

members in July 2010 of the NC and Western NC Conferences of the United Methodist Church (UMC). These two governing bodies employ roughly 1,800 clergy. We invited individuals based on clergy occupation status rather than health status. There were intentionally few exclusion criteria (e.g., being on leave, seminary professors, hospital chaplains).

Of the 1,745 clergy invited to participate in the trial, 1,114 (64%) clergy consented and provided survey and in-person cardiometabolic screening data in fall 2010. We randomly assigned these participants to three cohorts: the immediate intervention cohort (Cohort 1), a one-year waitlist intervention cohort (Cohort 2), and a two-year waitlist intervention cohort (Cohort 3). Only 49 participants received the intervention in a cohort different from their random assignment. Each cohort received two years of intervention, with Cohort 1 starting in 2011, Cohort 2 in 2012, and Cohort 3 in 2013. The timing enabled Cohort 3 to serve as a waitlist control group to Cohort 1 for the two-year outcomes paper.

We conducted cardiometabolic screenings with each of the cohorts immediately prior to intervention (0 months, called “baseline” in this paper) and at 12, 18, and 24 months of intervention. Cohorts 1 and 3 had additional measurements at 30, 36, and 42 months (i.e., 6, 12, and 18 months after intervention completion). Post-intervention data were not collected from Cohort 2. Although study blinding was not possible (participants knew when they received the intervention), the cardiometabolic screening assessors were blinded to cohort.

Duke University’s Arts & Sciences Institutional Review Board (IRB) approved all procedures. Study participants gave free and informed consent.

Measures

Staff measured height, weight, high-density lipoproteins, and triglycerides, as well as systolic and diastolic blood pressure, using detailed protocols (available upon request). We measured height to the nearest 0.25 inch using a Seca 213 stadiometer, and weight to the nearest 0.2 lb. on a calibrated digital scale (Seca 876), with the participant removing heavy clothing and shoes prior to being weighed. We created Body Mass Index (BMI) categories using the National Heart, Lung, and Blood Institute definition (NHLBI Expert Panel, 1998) [52].

We collected blood samples via finger sticks for lipid tests. High-density lipoproteins and triglycerides values were measured by the Cholestech LDX system. We asked participants to fast for at least 9 hours prior to data collection. After a five-minute resting period, blood pressure was assessed three times on the right arm using an Omron HEM-907XL machine, with a thirty-second rest between measurements. Throughout, participants remained seated with feet flat on the floor and arm resting at heart level. We took means of the three resting systolic and diastolic blood pressure values, respectively, for analysis. Staff recorded cardiometabolic data on paper then double-entered the data. We performed consistency checks across time periods to create a reliable final database.

Analyses

In the current analyses, we used the cohort in which participants ultimately received the intervention, allowing the analyses to accurately reflect the number of elapsed months since the end of the intervention. With our focus on post-intervention outcomes, the current analysis could only use: 1) data immediately before intervention and during intervention from all individuals and 2) post-intervention data from individuals who received the intervention with Cohorts 1 and 3 [39]. These post-intervention data were then referenced against baseline data

immediately prior to intervention receipt. There were a total of 719 participants who received the intervention in Cohorts 1 and 3. (See Supplement for sample size details.)

Health outcomes of interest, all continuous, were weight, percentage body weight change, high-density lipoproteins (for the entire sample and separately for males and females given different recommended levels by gender), systolic blood pressure, diastolic blood pressure, and triglycerides. We fitted a piecewise random effects linear regression model separately to each health outcome by regressing the outcome on three pieces of intervention (the first year in intervention, the second year in intervention, and the 18 months post-intervention) and on cohort (a 2-level factor coded as a variable indicating Cohort 1, with Cohort 3 as the reference level to allow for possible period effects due to different intervention starting time points). To account for correlation over time due to repeated measurements on the same individual, we included correlated random intercepts and random slopes which allowed for decreasing correlation of measurements further apart in time. Different random slopes were specified for each of the three time periods of the model to mirror the three-piece fixed effects model for time. We specified residual errors that were independent and normally distributed. Because triglycerides were both left-censored due to a lower limit of detection and right-censored due to a higher limit of detection, we fitted a Tobit piecewise random effects model for this outcome, in which three pieces of intervention and correlated random intercepts were included. We analyzed outcomes in the entire sample and in stratified samples by baseline BMI category (<25; 25 to <30; 30 to <40; and ≥ 40 kg/m²).

For each health outcome, we reported model-based estimates of the rate of change per year over each of the 3 pieces of intervention. We also obtained model-based estimates of mean change for each outcome from baseline to 18 months post-intervention (0-42 months), given our interest in determining whether weight loss can be maintained long-term. Standardized effect sizes for the primary outcome (weight) were obtained by scaling the mean weight changes (and their 95% CI) by the model-estimated residual standard deviation. All analyses were performed using Stata v15.1 software [53].

As in all follow-up studies, some participants missed data at some measurement time points, either because they dropped out of the study or because they were unable to attend the measurement visit. To ascertain whether the participants who had missing outcome data at a follow-up time point were different from those who were not missing at the same time point, we compared their baseline characteristics. Cohort, age, marital status, and education level were predictive of missing the primary outcome (weight) at follow-up; baseline weight and very low triglycerides values (≤ 45 mg/dL) at baseline had a trend of being predictive of missing weight at follow-up. In a sensitivity analysis to account for these identified predictors of missing outcome data, we adjusted for these covariates in our models of weight and percentage body weight change and then compared the results to our primary results; there were no substantive differences. Because the random effects linear regression modeling approach we used is likelihood-based, under an assumption of outcome data being missing-at-random (MAR) conditional on the identified covariates, the results from our sensitivity analyses provide valid estimates of changes over time, adjusting for the missing data pattern.

In another sensitivity analysis, participants who received the intervention with Cohort 2 were included in order to provide additional information on outcome levels during the intervention, although they provide no information on post-intervention outcome levels. This

sensitivity analysis found that the inclusion of data from Cohort 2 did not substantially alter the trends in findings.

The trial did not include a control group for time points beyond two years; we therefore wondered how the weight changes seen in Spirited Life participants would have compared to weight changes observed in those clergy who declined to enroll in the Spirited Life trial. In supportive analyses, we used data from a longitudinal panel survey conducted by our team with the same NC UMC clergy population. We were able to compare self-reported weight at two and four years following the start of the Spirited Life intervention between clergy who did and did not enroll in the trial. We elaborate the methods and results in the Supplement.

Results

Spirited Life participants followed post-intervention (n=719, i.e., Cohorts 1 and 3) were primarily male (69.0%), white (89.6%), and married (90.1%), with a mean age of 52.7 (SD=10.0 years) (see Table 1). All participants were employed by the UMC at the time of enrollment; at the start of the intervention, 98.3% reported having health insurance.

Table 2 displays the model-based estimated results of changes in outcomes: rate of change per year for each of the periods of time described in the Analyses, as well as overall change throughout the study period. At 12 months, change in mean weight was -2.4 kg (95% CI: -2.8 kg, -2.1 kg), corresponding to a standardized effect size of -0.75 (95% CI: -0.86, -0.65), or -2.4% (95% CI: -2.7%, -2.1%) of baseline body weight. For all participants combined, these improvements in weight slowly decayed, with a mean weight gain of 0.7 kg (95% CI: 0.3 kg, 1.0 kg) per year during the 18 months post-intervention (i.e., months 24-42). At the 42-month follow-up, mean weight did not significantly differ from baseline at -0.5 kg (95% CI: -1.2 kg, -0.2 kg), corresponding to a standardized effect size of -0.15 (-0.36, 0.06).

As shown in Table 2 and Figure 2, weight loss maintenance differed by baseline BMI status. Normal-weight participants with baseline BMI values <25 kg/m² experienced a mean weight gain of 1.7 kg (95% CI: 0.7 kg, 2.8 kg) from baseline to 42 months follow-up. Participants with baseline BMI values of 25 to <30 kg/m² (n=247) experienced a weight decrease at 12 months and gradually returned to their baseline weight at 42 months (0.2 kg, 95% CI: -0.9 kg, 1.3 kg). The same pattern of initial weight loss with returning to – but not exceeding – baseline levels was true for participants with baseline BMI values of 30 to <40 kg/m² (n=264). However, participants with baseline BMI values 40 kg/m² and higher (n=85) experienced greater weight loss at 12 months (-4.6 kg, 95% CI: -6.2 kg, -3.0 kg) relative to the other BMI groups, displayed non-significant weight gain between 12 and 24 months (0.2 kg, 95% CI: -2.0 kg, 2.4 kg) and non-significant weight loss post-intervention between 24 and 42 months (-0.9 kg per year, 95% CI: -2.7 kg, 0.9 kg). This high BMI group maintained their initial weight loss, with a mean loss of 5.8 kg (95% CI: -8.9 kg, -2.7 kg) from baseline to 42 months.

Additional to weight changes, triglycerides improved at 12 months (-14.6 mg/dL, 95% CI: -19.3 mg/dL, -10.0 mg/dL) and these improvements were sustained at 42 months (-13.9 mg/dL, 95% CI: -18.6 mg/dL, -9.2 mg/dL). Systolic and diastolic blood pressures (SBP; DBP) improved at 12 months (SBP: -3.8 mmHg, 95% CI: -4.6 mmHg, -2.9 mmHg; DBP: -2.0 mmHg, 95% CI: -2.6 mmHg, -1.4 mmHg). With some decay, mean SBP improvement remained significant at 42 months (-1.9 mmHg, 95% CI: -3.0 mmHg, -0.9 mmHg), although mean DBP returned to values

similar to baseline (-0.7 mmHg, 95% CI: -1.4 mmHg, 0.0 mmHg). For high-density lipoproteins, in men only, improvements occurring between 12 and 24 months (1.5 mg/dL, 95% CI: 0.8 mg/dL, 2.3 mg/dL) were partially sustained at 42 months (0.8 mg/dL compared to baseline, 95% CI: 0.0 mg/dL, 1.5 mg/dL). (See Supplement for these outcomes by BMI category.)

In supportive analyses, on average, compared to those who did not participate in Spirited Life, Spirited Life participants lost 1.9% (95% CI: 1.3%, 2.5%) more of their baseline body weight over 2 years, and lost 1.0% (95% CI: 0.1%, 1.8%) more of their baseline body weight over 4 years. This corresponds to -1.7 kg (95% CI: -2.3 kg, -1.1 kg) for Spirited Life participants compared to non-participants over 2 years and -1.1 kg (95% CI: -1.9 kg, -0.2 kg) over 4 years. Among clergy with BMI \geq 40 kg/m², Spirited Life participants lost 4.3 kg more over 2 years (95% CI: -7.8 kg, -0.9 kg) and 7.2 kg (95% CI: -13.4 kg, -1.0 kg) more over 4 years than non-participants.

Discussion

Participation in the Spirited Life intervention was consistent with achieving small short-term weight loss among clergy men and women in NC, and long-term weight loss maintenance among participants with BMI values of 40 kg/m² and above. This study aimed to test whether a lower intensity wellness intervention could be effective in reducing the prevalence of metabolic syndrome and obesity. It was not designed to produce large levels of weight loss. Yet, participation in the intervention's combination of Naturally Slim[®] online modules, health coaching, and social support was consistent with an average weight loss of 2.4 kg or 2.4% body weight loss at 12 months. Weight regains slowly occurred until 42 months from intervention start, notably without increases beyond pre-intervention levels, on average. These changes in weight were accompanied by small but clinically meaningful changes in cardiometabolic indicators, namely blood pressure and triglycerides, even at 42 months. In additional analyses, we were able to show that control-based comparisons support our claim that on average, small weight loss with partial weight loss maintenance at two years was attributable to the holistic Spirited Life intervention, and with complete weight loss maintenance at four years for those with BMI \geq 40 kg/m².

Obesity is typically recalcitrant to treatment and many individuals return to their pre-treatment weight. Without treatment, they may have started on a trajectory towards continued weight gain, as is common in adults. Although weight loss should always be the frontline treatment for individuals with obesity, weight stability for overweight and obese individuals not interested in weight loss is also a goal [6]. Interventions such as Spirited Life that slow progression towards inevitable weight gain offer health benefits.

An important finding of this study was different results by baseline BMI status. Unfortunately, participants with normal-weight BMI values at baseline did gain weight by 42 months. However, participants with BMI values \geq 40 kg/m² continued to lose weight post-intervention. It is possible that these participants with a greater weight problem sought to motivate themselves for the long-term and were able to enact the intentionally easy-to-remember weight loss strategies beyond the intervention.

The long-term follow-up in this study provides evidence that these findings are robust and adds to the literature. Only a few weight loss interventions have data on efficacy past 24

months. The Look AHEAD Study – considered one of the gold standard behavioral weight control trials – found that the intervention was effective in producing a 3% weight loss almost 10 years post-baseline [54]. Yet, this was arguably one of the most intensive interventions tested for producing weight loss. As such, it is not clear whether behavioral interventions are effective long-term, particularly when tested with a lower intensity approach.

The average decrease in systolic blood pressure at 42 months was 1.9 mmHg. Reductions in blood pressure following various pharmacologic and multi-component interventions significantly reduce the risk of future cardiovascular disease and mortality [55-57]. However, the mean reduction observed here is smaller than those established as conferring significant risk reduction, suggesting that only a subset of the Spirited Life participants with larger reductions would likely experience clinically meaningful blood pressure reductions. The decrease in triglycerides at 42 months was 13.9 mg/dL. Although improved lipid profiles similarly confer significant health benefits, the specific effects of reductions in triglycerides is currently not known [58]. However, the sustained nature of the improvements in blood pressure and triglycerides would increase the likelihood of health benefits.

Unlike many weight loss trials, the Spirited Life trial enrolled from a group of individuals who interact two or more times per year and who share an identity and affinity for each other. It is possible that even infrequent interaction allowed participants to support each other in weight loss and maintenance. Promoting health among respected leaders such as clergy may impact larger circles, as well. For example, clergy may not only influence each other, but may also influence the health of their congregants [59].

The strengths of this study include having 3.5 years of data and measured weight and cardiometabolic indicators for a large number of participants. In addition, this intervention study enrolled a large number of men (69%), who tend to be under-represented in controlled weight trials [60]. The limited trials with men have focused on inducing weight loss, whereas the current study additionally examined weight loss maintenance in men [61; 62]. The current study was also unique in using a wellness strategy that included mental and physical health and spiritual well-being. This study's limitations include potential lack of generalizability due to its predominantly white, religious sample. In addition, the trial itself did not include a control group for time points beyond two years. However, we used other data we had on members of the population who declined study participation, to offer findings on weight changes over time in the absence of the Spirited Life intervention. Admittedly, this is not an ideal control group and those who declined study participation may simply be unconcerned about weight.

In conclusion, we found that participation in this holistic wellness intervention, Spirited Life, was consistent with prevention of weight gain for participants with BMI values between 25 and 40 kg/m², and complete weight loss maintenance for participants with BMI values ≥40 kg/m², at 42 months post-baseline. Regardless of BMI status, Spirited Life participation was also consistent with small but clinically meaningful changes in cardiometabolic indicators at 42 months post-baseline. Weight loss maintenance programs should consider lower-intensity, two-year interventions.

This study was funded by a grant from the Rural Church Area of The Duke Endowment. Rae Jean Proeschold-Bell, Dori Steinberg, Jia Yao, David E. Eagle, Timothy W. Smith, Grace Cai, and Elizabeth L. Turner declare that they have no conflicts of interest. All procedures performed in

studies involving human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. The study protocol was approved by the Duke University Arts & Science Institutional Review Board and the Westat Institutional Review Board. Informed consent was obtained from all individual participants included in the study. This article does not contain any studies with animals performed by any of the authors.

Acknowledgments

This trial was made possible by the United Methodist clergy of North Carolina, along with support from their bishops and leaders. David Toole served as principal investigator and Carl Weisner as senior director. Data were collected and organized by Crystal MacAllum, Gail Thomas, Ed Mann, and their team at Westat, as well as Rachel Meyer, Amanda Wallace, Xiang-Fang Li, Alana Bennett, Melanie Kolkin, and Holly Hough. Key intervention decisions were made by Gary G. Bennett, Robin Swift, and H. Edgar Moore, with Redford Williams and Virginia Williams supporting the Williams LifeSkills component. Thirteen Wellness Advocates implemented the intervention with great care. We thank Celia Hybels, Glen Milstein, Jennifer Lutz, Carl Weisner, and Rachel Meyer for their insightful comments during the writing process. This trial was funded by the Rural Church Area of The Duke Endowment.

References

1. Haslam, D. W., & James, W. P. T. (2005). Obesity. *The Lancet*, 366(9492), 1197-1209.
2. Poobalan, A. S., Aucott, L. S., Smith, W. C. S., Avenell, A., Jung, R., & Broom, J. (2007). Long-term weight loss effects on all cause mortality in overweight/obese populations. *Obesity Reviews*, 8(6), 503-513.
3. Unick, J. L., Beavers, D., Bond, D. S., Clark, J. M., Jakicic, J. M., Kitabchi, A. E., . . . Wing, R. R. (2013). The long-term effectiveness of a lifestyle intervention in severely obese individuals. *The American Journal of Medicine*, 126(3), 236-242. e232.
4. Look AHEAD Research Group. (2007). Reduction in weight and cardiovascular disease risk factors in individuals with type 2 diabetes: One-year results of the Look AHEAD Trial. *Diabetes Care*, 30(6), 1374.
5. Wing, R. R. (2004). Behavioral approaches to the treatment of obesity. *Handbook of obesity: Clinical applications*, 2, 147-167.
6. Jensen, M. D., Ryan, D. H., Apovian, C. M., Ard, J. D., Comuzzie, A. G., Donato, K. A., . . . Kushner, R. F. (2014). 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Journal of the American College of Cardiology*, 63(25 Part B).
7. Wing, R. R., & Phelan, S. (2005). Long-term weight loss maintenance. *The American Journal of Clinical Nutrition*, 82(1), 222S-225S.
8. Avenell, A., Broom, J., Brown, T. J., Poobalan, A., Aucott, L., Stearns, S., . . . Grant, A. M. (2004). Systematic review of the long-term effects and economic consequences of treatments for

obesity and implications for health improvement *NIHR Health Technology Assessment programme: Executive Summaries*. Southampton (UK): NIHR Journals Library.

9. Dombrowski, S. U., Avenell, A., & Sniehot, F. F. (2010). Behavioural interventions for obese adults with additional risk factors for morbidity: systematic review of effects on behaviour, weight and disease risk factors. *Obesity Facts*, 3(6), 377-396.
10. Svetkey, L. P., Stevens, V. J., Brantley, P. J., Appel, L. J., Hollis, J. F., Loria, C. M., . . . Smith, P. (2008). Comparison of strategies for sustaining weight loss: The weight loss maintenance randomized controlled trial. *JAMA*, 299(10), 1139-1148.
11. Mann, T., Tomiyama, A. J., Westling, E., Lew, A.-M., Samuels, B., & Chatman, J. (2007). Medicare's search for effective obesity treatments: Diets are not the answer. *American Psychologist*, 62(3), 220.
12. Jeffery, R. W., Epstein, L. H., Wilson, G. T., Drewnowski, A., Stunkard, A. J., & Wing, R. R. (2000). Long-term maintenance of weight loss: current status. *Health Psychology*, 19(1S), 5.
13. Wing, R. R. (2002). Behavioral weight control. *Handbook of obesity treatment*, 2, 301-317.
14. Wadden, T. A., Neiberg, R. H., Wing, R. R., Clark, J. M., Delahanty, L. M., Hill, J. O., . . . Vitolins, M. Z. (2011). Four-Year Weight Losses in the Look AHEAD Study: Factors Associated With Long-Term Success. *Obesity*, 19(10), 1987-1998.
15. Penn, L., White, M., Lindström, J., den Boer, A. T., Blaak, E., Eriksson, J. G., . . . Walker, M. (2013). Importance of weight loss maintenance and risk prediction in the prevention of type 2 diabetes: Analysis of European Diabetes Prevention Study RCT. *PLoS One*, 8(2), e57143.
16. Zheng, Y., Manson, J. E., Yuan, C., Liang, M. H., Grodstein, F., Stampfer, M. J., . . . Hu, F. B. (2017). Associations of weight gain from early to middle adulthood with major health outcomes later in life. *JAMA*, 318(3), 255-269.
17. Mozaffarian, D., Hao, T., Rimm, E. B., Willett, W. C., & Hu, F. B. (2011). Changes in diet and lifestyle and long-term weight gain in women and men. *New England Journal of Medicine*, 364, 2392-2404.
18. Truesdale, K. P., Stevens, J., Lewis, C. E., Schreiner, P. J., Loria, C. M., & Cai, J. (2006). Changes in risk factors for cardiovascular disease by baseline weight status in young adults who maintain or gain weight over 15 years: The CARDIA study. *International Journal of Obesity*, 30(9), 1397-1407.
19. Montesi, L., El Ghoch, M., Brodosi, L., Calugi, S., Marchesini, G., & Dalle Grave, R. (2016). Long-term weight loss maintenance for obesity: a multidisciplinary approach. *Diabetes, metabolic syndrome and obesity: targets and therapy*, 9, 37.
20. MacLean, P. S., Wing, R. R., Davidson, T., Epstein, L., Goodpaster, B., Hall, K. D., . . . Rosenbaum, M. (2015). NIH working group report: Innovative research to improve maintenance of weight loss. *Obesity*, 23(1), 7-15.
21. Hill, J. O., Wyatt, H., Phelan, S., & Wing, R. (2005). The National Weight Control Registry: is it useful in helping deal with our obesity epidemic? *Journal of Nutrition Education and Behavior*, 37(4), 206-210.
22. Wing, R. R., Tate, D. F., Gorin, A. A., Raynor, H. A., & Fava, J. L. (2006). A self-regulation program for maintenance of weight loss. *New England Journal of Medicine*, 355(15), 1563-1571.
23. Riebe, D., Blissmer, B., Greene, G., Caldwell, M., Ruggiero, L., Stillwell, K. M., & Nigg, C. R. (2005). Long-term maintenance of exercise and healthy eating behaviors in overweight adults. *Preventive Medicine*, 40(6), 769-778.

24. Heinen, L., & Darling, H. (2009). Addressing Obesity in the Workplace: The Role of Employers. *The Milbank Quarterly*, 87(1), 101-122.
25. Williams, A. E., Stevens, V. J., Albright, C. L., Nigg, C. R., Meenan, R. T., & Vogt, T. M. (2014). The results of a 2-year randomized trial of a worksite weight management intervention. *American Journal of Health Promotion*, 28(5), 336-339.
26. Østbye, T., Stroot, M., Eisenstein, E. L., & Dement, J. M. (2016). The effects of two workplace weight management programs and weight loss on health care utilization and costs. *Journal of Occupational and Environmental Medicine*, 58(2), 162.
27. Wilcox, A. (2010). Hospital system cohort case study [PowerPoint presentation]. Durham, NC: ACAP Health.
28. Miller, W. R., & Rollnick, S. (1991). *Motivational interviewing: Preparing people to change addictive behavior*. New York: Guilford Press.
29. Van Dorsten, B. (2007). The use of motivational interviewing in weight loss. *Current Diabetes Reports*, 7(5), 386-390.
30. Elley, C. R., Kerse, N., Arroll, B., & Robinson, E. (2003). Effectiveness of counselling patients on physical activity in general practice: cluster randomised controlled trial. *BMJ*, 326(7393), 793.
31. Burke, B. L., Arkowitz, H., & Menchola, M. (2003). The efficacy of motivational interviewing: a meta-analysis of controlled clinical trials. *Journal of Consulting and Clinical Psychology*, 71(5), 843.
32. Hettema, J., Steele, J., & Miller, W. R. (2005). Motivational interviewing. *Annual Review of Clinical Psychology*, 1, 91-111.
33. Appel, L. J., Clark, J. M., Yeh, H.-C., Wang, N.-Y., Coughlin, J. W., Daumit, G., . . . Geller, S. (2011). Comparative effectiveness of weight-loss interventions in clinical practice. *New England Journal of Medicine*, 365(21), 1959-1968.
34. Butterworth, S. W., Linden, A., & McClay, W. (2007). Health coaching as an intervention in health management programs. *Disease Management & Health Outcomes*, 15(5), 299-307.
35. Butterworth, S., Linden, A., McClay, W., & Leo, M. C. (2006). Effect of motivational interviewing-based health coaching on employees' physical and mental health status. *Journal of Occupational Health Psychology*, 11(4), 358.
36. Linden, A., Butterworth, S. W., & Prochaska, J. O. (2010). Motivational interviewing-based health coaching as a chronic care intervention. *Journal of Evaluation in Clinical Practice*, 16(1), 166-174.
37. Vale, M. J., Jelinek, M. V., Best, J. D., Dart, A. M., Grigg, L. E., Hare, D. L., . . . McNeil, J. J. (2003). Coaching patients On Achieving Cardiovascular Health (COACH): A Multicenter Randomized Trial in Patients With Coronary Heart Disease. *Archives of Internal Medicine*, 163(22), 2775-2783.
38. Svetkey, L. P., Batch, B. C., Lin, P. H., Intille, S. S., Corsino, L., Tyson, C. C., . . . Bennett, G. G. (2015). Cell phone intervention for you (CITY): A randomized, controlled trial of behavioral weight loss intervention for young adults using mobile technology. *Obesity (Silver Spring)*, 23(11), 2133-2141.
39. Proeschold-Bell, R. J., Swift, R., Moore, H. E., Bennett, G., Li, X. F., Blouin, R., . . . Toole, D. (2013). Use of a randomized multiple baseline design: Rationale and design of the Spirited Life holistic health intervention study. *Contemporary Clinical Trials*, 35(2), 138-152.
doi:10.1016/j.cct.2013.05.005

40. Ries, N. M. (2012). Financial Incentives for Weight Loss and Healthy Behaviours. *Healthcare Policy*, 7(3), 23-28.
41. Yancy Jr, W. S., Shaw, P. A., Wesby, L., Hilbert, V., Yang, L., Zhu, J., . . . Volpp, K. G. (2018). Financial incentive strategies for maintenance of weight loss: results from an internet-based randomized controlled trial. *Nutrition & Diabetes*, 8(1), 33. doi:10.1038/s41387-018-0036-y
42. Institute of Medicine. (2014). Institute of Medicine classifications for prevention. Retrieved from http://dpbh.nv.gov/uploadedFiles/mhnavgov/content/Meetings/Bidders_Conference/Institute%20of%20Medicine%20Prevention%20Classifications-rev10.20.14.pdf
43. Proeschold-Bell, R. J., & LeGrand, S. (2010). High rates of obesity and chronic disease among United Methodist clergy. *Obesity*, 18(9), 1867-1870.
44. Halaas, G. W. (2002). *Ministerial health and wellness: 2002 Evangelical Lutheran Church in America* (Division for Ministry, Board of Pensions Ed.). Chicago, IL Evangelical Lutheran Church in America.
45. Center for Health, Wespeth Benefits and Investments. (2017). Well-being survey of the United Methodist Church active U.S. clergy—2017. Retrieved from <https://www.wespeth.org/assets/1/7/5058.pdf>
46. Lindholm, G., Johnston, J., Dong, F., Moore, K., & Ablah, E. (2016). Clergy wellness: An assessment of perceived barriers to achieving healthier lifestyles. *Journal of Religion and Health*, 55(1), 97-109.
47. Baruth, M., Wilcox, S., & Evans, R. (2014). The health and health behaviors of a sample of African American pastors. *Journal of Health Care for the Poor and Underserved*, 25(1), 229-241.
48. Webb, B., Bopp, M., & Fallon, E. (2013). Factors associated with obesity and health behaviors among clergy. *Health Behaviour & Public Health*, 3(1), 20-28.
49. Proeschold-Bell, R. J., Turner, E. L., Bennett, G. G., Yao, J., Li, X.-F., Eagle, D. E., . . . Toole, D. C. (2017). A 2-year holistic health and stress intervention: Results of an RCT in clergy. *American Journal of Preventive Medicine*, 53(3), 290-299. doi:10.1016/j.ampere.2017.04.009
50. Alberti, K. G. M. M., Zimmet, P., & Shaw, J. (2006). Metabolic syndrome—a new world-wide definition. A consensus statement from the International Diabetes Federation. *Diabetic Medicine*, 23(5), 469-480.
51. Dunkley, A. J., Bodicoat, D. H., Greaves, C. J., Russell, C., Yates, T., Davies, M. J., & Khunti, K. (2014). Diabetes prevention in the real world: effectiveness of pragmatic lifestyle interventions for the prevention of type 2 diabetes and of the impact of adherence to guideline recommendations: a systematic review and meta-analysis. *Diabetes Care*, 37(4), 922-933.
52. NHLBI Obesity Education Initiative Expert Panel on the Identification, Evaluation, and Treatment of Obesity in Adults (US). (1998). *Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: The evidence report*. Bethesda, MD: National Heart, Lung, and Blood Institutes.
53. StataCorp. (2017). *Stata Statistical Software: Release 15*. College Station, TX: StataCorp LLC.
54. Dutton, G. R., & Lewis, C. E. (2015). The Look AHEAD Trial: implications for lifestyle intervention in type 2 diabetes mellitus. *Progress in Cardiovascular Diseases*, 58(1), 69-75.
55. Bundy, J. D., Li, C., Stuchlik, P., Bu, X., Kelly, T. N., Mills, K. T., . . . He, J. (2017). Systolic blood pressure reduction and risk of cardiovascular disease and mortality: A systematic review and network meta-analysis. *JAMA cardiology*, 2(7), 775-781.

56. Ettehad, D., Emdin, C. A., Kiran, A., Anderson, S. G., Callender, T., Emberson, J., . . . Rahimi, K. (2016). Blood pressure lowering for prevention of cardiovascular disease and death: A systematic review and meta-analysis. *The Lancet*, *387*(10022), 957-967.
57. Sundström, J., Arima, H., Jackson, R., Turnbull, F., Rahimi, K., Chalmers, J., . . . Neal, B. (2015). Effects of blood pressure reduction in mild hypertension: A systematic review and meta-analysis. *Annals of Internal Medicine*, *162*(3), 184-191.
58. Nordestgaard, B. G. (2016). Triglyceride-rich lipoproteins and atherosclerotic cardiovascular disease: New insights from epidemiology, genetics, and biology. *Circulation Research*, *118*(4), 547-563.
59. Boyatzis, R. E., Brizz, T., & Godwin, L. N. (2011). The effect of religious leaders' emotional and social competencies on improving parish vibrancy. *Journal of Leadership and Organizational Studies*, *18*(2), 192-206.
60. Pagoto, S. L., Schneider, K. L., Oleski, J. L., Luciani, J. M., Bodenlos, J. S., & Whited, M. C. (2012). Male inclusion in randomized controlled trials of lifestyle weight loss interventions. *Obesity*, *20*(6), 1234-1239.
61. Crane, M. M., Ward, D. S., Lutes, L. D., Bowling, J. M., & Tate, D. F. (2016). Theoretical and behavioral mediators of a weight loss intervention for men. *Annals of Behavioral Medicine*, *50*(3), 460-470.
62. Morgan, P. J., Lubans, D. R., Collins, C. E., Warren, J. M., & Callister, R. (2009). The SHED-IT randomized controlled trial: Evaluation of an internet-based weight-loss program for men. *Obesity*, *17*(11), 2025-2032.

Table 1
Baseline characteristics of the study participants (N=719)

Characteristics	% (n/N) or M (SD), n
Gender, % (n/N)	
Male	69.0% (496/719)
Female	31.0% (223/719)
Age, M (SD), N	52.7 (10.0), 719
Race, % (n/N)	
White	89.6% (644/719)
Black	6.0% (43/719)
Other	4.5% (32/719)
Marital status ^a , % (n/N)	
Married	90.1% (647/718)
Not married	9.9% (71/718)
Education ^a , % (n/N)	
College & below	15.3% (110/718)
Master's degree	71.7% (515/718)
Doctoral degree	13.0% (93/718)
Health insurance ^a , % (n/N)	
Insured	98.3% (707/719)
Not insured	1.7% (12/719)
Financial stress ^a , % (n/N)	
High stress	17.8% (128/718)
Moderate or low	82.2% (590/718)
Weight in KG, M (SD), N	93.8 (23.3), 719
BMI categories, % (n/N)	
BMI <25	17.1% (123/719)
25 ≤ BMI <30	34.4% (247/719)
30 ≤ BMI <40	36.7% (264/719)
BMI ≥40	11.8% (85/719)
Metabolic syndrome, % (n/N)	34.7% (249/717)
High-density lipoproteins, M (SD), N	45.9 (15.4), 713
High-density lipoproteins in males	40.9 (12.5), 491
High-density lipoproteins in females	56.9 (15.5), 222
Systolic blood pressure, M (SD), N	124.3 (14.6), 717
Diastolic blood pressure, M (SD), N	77.8 (10.2), 717
Triglycerides	
Low triglycerides (<45), % (n/N)	3.2% (23/712)
Triglycerides values (≥45), M (SD), N	141.6 (82.6), 689/712
Intervention-based cohort, % (n/N)	
Cohort 1: Immediate intervention	49.5% (356/719)
Cohort 3: Two-year waitlist	50.5% (363/719)

Note. ^aMissing demographical values are filled using values from the two other years. This was true for marital status (n=23), education (n=8), health insurance (n=24), and financial stress (n=25).

Table 2

Results of modeling Spirited Life health outcomes during and after intervention

Weight outcomes	Percentage of baseline weight, all (N=719)	Weight, kg, all (N=719)	Weight, kg, BLBMI<25 (N=123)	Weight, kg, 25≤BL BMI<30 (N=247)	Weight, kg, 30≤BL BMI<40 (N=264)	Weight, kg, BL BMI≥40 (N=85)
Rate of change in outcome per year						
0-12 mths (int)	-2.40 [-2.73, -2.08]; p<.001	-2.40 [-2.75, -2.06]; p<.001	-0.07 [-0.51, 0.37]; p=.749	-2.12 [-2.55, -1.69]; p<.001	-3.05 [-3.64, -2.45]; p<.001	-4.59 [-6.23, -2.95]; p<.001
12-24 mths (int)	1.16 [0.72, 1.60]; p<.001	0.93 [0.53, 1.34]; p<.001	0.92 [0.41, 1.44]; p<.001	0.98 [0.48, 1.47]; p<.001	1.18 [0.52, 1.84]; p<.001	0.19 [-1.98, 2.36]; p=.865
24-42 mths (post int)	0.93 [0.43, 1.43]; p<.001	0.67 [0.31, 1.02]; p<.001	0.57 [0.11, 1.04]; p=.016	0.91 [0.35, 1.46]; p=.001	1.00 [0.42, 1.58]; p<.001	-0.91 [-2.74, 0.92]; p=.329
Overall change						
0-42 mths	0.15 [-0.93, 1.22]; p=.788	-0.47 [-1.15, 0.20]; p=.171	1.71 [0.66, 2.75]; p=.001	0.22 [-0.88, 1.31]; p=.695	-0.36 [-1.47, 0.75]; p=.525	-5.76 [-8.85, -2.67]; p<.001
Other health outcomes						
Other health outcomes	HDL, mg/dL, all (N=719)	HDL, mg/dL, males (N=496)	HDL, mg/dL, females (N=223)	Systolic blood pressure, mmHg (N=719)	Diastolic blood pressure, mmHg (N=719)	Triglycerides, mg/dL (N=713)
Rate of change in outcome per year						
0-12 mths (int)	-0.86 [-1.49, -0.23]; p=.008	-0.45 [-1.16, 0.25]; p=.207	-1.80 [-3.09, -0.51]; p=.006	-3.75 [-4.63, -2.87]; p<.001	-2.01 [-2.59, -1.44]; p<.001	-14.64 [-19.28, -10.01]; p<.001
12-24 mths (int)	2.21 [1.50, 2.92]; p<.001	1.54 [0.79, 2.28]; p<.001	3.68 [2.13, 5.22]; p<.001	1.82 [0.90, 2.75]; p<.001	0.51 [-0.12, 1.13]; p=.113	2.36 [-2.36, 7.08]; p=.327
24-42 mths (post int)	-0.55 [-1.00, -0.10]; p=.016	-0.22 [-0.72, 0.29]; p=.400	-1.29 [-2.21, -0.38]; p=.006	-0.01 [-0.71, 0.70]; p=.985	0.56 [0.11, 1.01]; p=.016	-1.09 [-4.37, 2.18]; p=.513
Overall change						
0-42 mths	0.52 [-0.20, 1.25]; p=.157	0.76 [0.00, 1.52]; p=.049	-0.06 [-1.64, 1.53]; p=.942	-1.94 [-2.99, -0.88]; p<.001	-0.67 [-1.38, 0.04]; p=.065	-13.92 [-18.62, -9.22]; p<.001

Note. Independent variables are intervention levels in three pieces and the cohort dummy variable. Coefficient, 95% confidence interval and p value are reported. Overall changes are estimated using linear combinations of the model parameters.

BL = baseline

int = intervention

HDL = high-density lipoproteins

		Cohort 1		Cohort 2		Cohort 3	
		Data	Intervention	Data	Intervention	Data	Intervention
2010	Nov						
	Dec						
2011	Jan		3-day workshop including theological and stress management content		Jan 2012–Dec 2013 Cohort 2’s services mirror those of Cohort 1, with small improvements to smooth out delivery; these included more choices for when to start Naturally Slim, although all start times were in 2012.		Jan 2013–Dec 2014 Cohort 3’s services mirror those of Cohort 2, with a change to online stress management content that participants generally did not engage with.
	Feb						
	Mar		10-week online weight loss program; 1st health coaching call				
	Apr						
	May						
	June		Optional monthly health coaching calls				
	July						
	Aug						
	Sept		Health coaching on stress mgmt. skills				
	Oct		2-day workshop with theological and process of health behavior change content; monthly health coaching calls				
	Nov		Health coaching call				
	Dec						
2012	Jan		\$500 grant & health goal setting; monthly health coaching calls				
	Feb						
	Mar		Monthly health coaching calls				
	Apr						
	May						
	June						
	July						
	Aug						
	Sept						
	Oct		1-day workshop; monthly health coaching calls				
	Nov						
	Dec						
2013	Jan						
	Feb						
	Mar						
	Apr						
	May						
	June						
	July						
	Aug						
	Sept						
	Oct						
	Nov						
	Dec						
2014	Jan						
	Feb						
	Mar						
	Apr						
	May						
	June						
	July						
	Aug						
	Sept						
	Oct						
	Nov						
	Dec						

Figure 1. Spirited Life intervention components by month and cohort. *Note.* “Data” indicates data collection which always included a survey and an in-person cardiometabolic screening.




Figure 2. Estimated weight change during and after the Spirited Life intervention by BMI category. BMI = Body Mass Index measured at baseline